



University of Hamburg, Jungiusstr. 11, 20355 Hamburg

Press release

University of Hamburg

Jungiusstr. 9A-11A
20355 Hamburg

Heiko Fuchs
Public Relation

Tel.: +49 - 40 - 428 38 - 69 59
Fax: +49 - 40 - 428 38 - 24 09
hfuchs@physnet.uni-hamburg.de

Hamburg, 2015/09/07

Realization of the building block of a Hund's metal

As reported in a recent publication of the journal **Nature Nanotechnology** a team of experimentalists and theoreticians of the University of Hamburg in cooperation with the University of Bremen, the Radboud University in Nijmegen and the Institute of Physics of the Czech Academy of Sciences have experimentally realized and studied the basic building block of an electron phase named **Hund's metal** which is present in some of the high-temperature superconductors.

The electronic properties of solid state materials used in today's electronic devices are governed by the properties of the electrons in their basic constituents, the atoms. As already realized almost ninety years ago by German physicist Friedrich Hund (1896 - 1997) the occupation of the atomic orbitals with electrons (see Figure), which is governed by their mutual repulsion due to their negative charge, leads to a peculiar ordering of the *spin* of these electrons which, in layman's terms, is the sense of rotation of the spinning motion of the electrons: the electrons all tend to spin with the same sense of rotation, a rule of thumb which is called *Hund's rule*.

Since the electron current in electronic devices consists precisely of these electrons hopping from one atom to the neighboring atom, the consolidation of the spinning motion of electrons due to Hund's rule may have profound consequences for the electronic properties of the device. Metallic materials in which the electron motion is governed by Hund's rule are called Hund's metals. Indeed, theorists have argued that the electrons in a recently heavily studied class of *superconductors* behave like Hund's metals. In these materials, the electrons are hopping without any resistance and thus can flow through the material without any loss of energy. So far, the superconducting electron flow does only survive at very low temperatures in these materials, and the corresponding devices therefore need to be cooled down to temperatures which are usually present in outer space only. The research community is therefore frantically searching for new materials, hopefully showing superconductivity under ambient conditions, which would solve some of the most pressing problems of the current era of information technology. However, for a target-oriented search of such materials, the electron properties of the basic constituents of Hund's metals need to be understood in detail, and this requirement was lacking so far.

A team of experimentalists and theoreticians of the University of Hamburg in cooperation with the University of Bremen, the Radboud University in Nijmegen and the Institute of Physics of the Czech Academy of Sciences have now experimentally realized such a basic constituent, which they coined *Hund's impurity*, by depositing iron-hydrogen-molecules on the surface of platinum (see Figure). Moreover, they were able to intentionally remove hydrogen from such a Hund's impurity by using the tip of a scanning tunneling microscope as a tool. The team found that attaching or removing the hydrogen has profound consequences for the electronic properties of the Hund's impurity which they studied in great detail by comparing the experimental data to cutting edge computer simulations of the system. In a next step, the researchers hope to be able to couple many Hund's impurities by moving them closer, again by using the tip of a scanning tunneling microscope as a tool. This would enable a bottom-up assembly of a Hund's metal and its study hopefully will give relevant insight for the targeted development of novel high-temperature superconducting materials.




Figure:

Left panel: Occupation of five electron orbitals (boxes) of an atom with five or six spin up (magenta colored arrows) or spin down (cyan colored arrows) electrons according to Hund's rules. For adding the sixth electron to the orbitals the energy of U_{Coulomb} has to be paid due to the mutual electrostatic repulsion of the negatively charged electrons. However, if one of the electrons changes its spin from up to down, an energy also has to be paid (J_{Hund}).

Center panel: Scanning tunneling microscope image of an iron atom (cone with red tip) and three iron-hydrogen-molecules (cones with yellow tips) on the surface of platinum.

Right panel: The hydrogen of the bottom right iron-hydrogen-molecule has been removed by using the tip of the scanning tunneling microscope as a tool.

Original Publication:

Tuning emergent magnetism in a Hund's impurity,

A. A. Khajetoorians, M. Valentyuk, M. Steinbrecher, T. Schlenk, A. Shick, J. Kolorenc, A. I. Lichtenstein, T. O. Wehling, R. Wiesendanger and J. Wiebe

Nature Nanotechnology (2015)

DOI: 10.1038/nnano.2015.193

Research funding:

EU: ERC Advanced Grant ASTONISH (Principal Investigator: Prof. Dr. Roland Wiesendanger)

DFG: SFB 668

Additional web sites:

<http://www.nanoscience.de>

<http://www.sfb668.de>

<http://www.nanoscience.de/astonish>

Additional information:

Dr. Jens Wiebe and
Dipl.-Chem. Heiko Fuchs
Universität Hamburg
Jungiusstr. 9A
20355 Hamburg

Tel.: +49- 40 - 4 28 38 - 32 82

Tel.: +49- 40 - 4 28 38 - 69 59

Fax.: +49- 40 - 4 28 38 - 24 09

E-Mail: jwiebe@physnet.uni-hamburg.de

E-Mail: hfuchs@physnet.uni-hamburg.de